Reference

Nuclides Analysis Result of the Radioactive Materials in the Air at the Opening of Buildings at Fukushima Daiichi NPS

(Data summarized on June 26)

Place of Sampling	Exhaust Facility of Granular Solid Strage (Outlet)						Density Limit Specified by the Reactor Regulation
Time of Sampling	Jun 20, 2013 2:30 PM - 2:40 PM						
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm ³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm ³)	Scaling Factor (①/②)	is specified in section 4 of Appendix 2)
I-131 (Approx. 8 days)	ND	1					1E-03
Cs-134 (Approx. 2 years)	ND	-					2E-03
Cs-137 (Approx. 30 years)	ND	-					3E-03

^{*} The radioactivity density is the sum of the volatile nuclides density and the particulate nuclides density.

O.OE-O is the same as $O.O \times 10^{-O}$

Data of other nuclides is under examination.

The detection limits are as follows.

Volatile; I-131: Approx. 2E-6Bq/cm³, Cs-134: Approx. 6E-6Bq/cm³, Cs-137: Approx. 8E-6Bq/cm³

Particulate; I-131: Approx. 1E-6Bq/cm³, Cs-134: Approx. 3E-6Bq/cm³, Cs-137: Approx. 4E-6Bq/cm³

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

 $[\]ensuremath{^{*}}$ "ND" indicates that the measurement result is below the detection limit.