

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Northern dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.7)	ND (3.8)	ND
Common skate (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.6)	6.5	6.5
Sea bass (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.5)	ND (4.3)	ND
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.6)	6.7	6.7
Ovalipes punctatus (whole)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.4)	ND (3.1)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (3.7)	ND (3.9)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2016	ND (4.2)	3.9	3.9
Northern dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.4)	ND (4.0)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (4.0)	ND (3.4)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (4.2)	ND (3.8)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lepidotrigla microptera (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.9)	ND (3.9)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.0)	3.7	3.7
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.0)	5.0	5.0
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (4.5)	ND (4.3)	ND
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (4.1)	ND (3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.6)	8.3	8.3
Pacific cod (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (3.2)	ND (3.6)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2016	ND (2.5)	ND (3.7)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (3.3)	ND (4.1)	ND
Lepidotrigla microptera (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (3.4)	3.7	3.7

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (3.6)	8.7	8.7
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (4.7)	6.8	6.8
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (3.4)	ND (3.5)	ND
Pacific cod (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2016	ND (5.0)	ND (4.2)	ND
Northern dogfish (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.1)	ND (3.6)	ND
Stone flounder (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.6)	6.7	6.7
Blue crab (whole)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.9)	ND (4.3)	ND
Lepidotrigla microptera (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.8)	ND (3.6)	ND
Schlegel's black rockfish (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.3)	ND (4.3)	ND
Sea raven (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (4.2)	3.9	3.9

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (4.8)	7.0	7.0
Microstomus achne (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	9.7	49	58.7
Flatfish (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (4.2)	8.9	8.9
Sea robin (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.6)	ND (3.3)	ND
Littlemouth flounder (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (4.2)	ND (3.8)	ND
Marbled sole (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (4.1)	4.9	4.9
Roundnose flounder (muscle)	Aruond 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 11, 2016	ND (3.4)	ND (3.2)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (4.1)	19	19
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (4.3)	26	26
Sebastes cheni (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (4.3)	21	21

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Drumfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (4.0)	ND (4.1)	ND
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (3.5)	10	10
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (4.2)	3.5	3.5
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 19, 2016	ND (3.8)	15	15
Greenling (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (4.0)	4.8	4.8
Northern dogfish (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (4.1)	ND (4.2)	ND
Schlegel's black rockfish (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (3.8)	3.7	3.7
Common skate (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	5.6	29	34.6
Microstomus achne (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (3.7)	12	12
Flatfish (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (3.5)	ND (4.0)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Dasyatis matsubarai (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	ND (4.5)	ND (3.1)	ND
Spotbelly rockfish (muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	May 19, 2016	9.0	54	63
Northern dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.3)	ND (3.7)	ND
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.6)	ND (3.6)	ND
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.6)	ND (3.9)	ND
Lepidotrigla microptera (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (4.2)	ND (3.9)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (4.0)	ND (4.5)	ND
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (4.5)	20	20
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.2)	ND (3.4)	ND
Dasyatis matsubarai (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (4.1)	ND (3.3)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (4.6)	3.5	3.5
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.7)	ND (3.7)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2016	ND (3.9)	ND (4.0)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.3)	ND (3.8)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.5)	ND (3.5)	ND
Lophius litilon (whole)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (4.5)	ND (4.1)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.4)	9.0	9.0
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (4.7)	ND (3.9)	ND
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.5)	ND (4.2)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.4)	ND (4.4)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 20, 2016	ND (3.6)	ND (3.7)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.3)	4.2	4.2
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (4.2)	ND (4.5)	ND
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (4.0)	ND (3.1)	ND
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.7)	6.9	6.9
Shark-skin flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.8)	ND (3.3)	ND
Loliginid (whole)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.9)	ND (4.4)	ND
Pointhead flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.1)	ND (4.1)	ND
Cloudy catshark (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.1)	ND (3.5)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.6)	ND (3.9)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (4.1)	4.4	4.4
Korean flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.2)	ND (3.5)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (4.8)	ND (4.3)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.8)	ND (4.4)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (2.3)	ND (4.1)	ND
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.8)	ND (3.5)	ND
Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.9)	ND (4.4)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.8)	ND (3.7)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.5)	ND (3.8)	ND
Slippery sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 20, 2016	ND (3.4)	ND (3.2)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (4.3)	ND (4.4)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.3)	ND (3.5)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.2)	ND (3.9)	ND
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (4.3)	11	11
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.8)	9.1	9.1
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (4.1)	ND (3.8)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.5)	ND (3.7)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.8)	3.6	3.6
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (3.6)	3.9	3.9
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	May 24, 2016	ND (4.4)	ND (3.4)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding inside the port)

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lepidotrigla microptera (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.9)	ND (3.2)	ND
Lophius litilon (whole)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.1)	ND (3.5)	ND
Common skete (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (4.1)	6.0	6.0
Microstomus achne (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.7)	5.0	5.0
Flatfish (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (4.2)	ND (3.7)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.5)	ND (3.9)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.7)	3.6	3.6
Marbled sole (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (4.3)	ND (3.5)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Daini NPS (T-B4)	May 24, 2016	ND (3.2)	ND (4.0)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, shown as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.