


Sampling points of drainage water at Fukushima Daiichi Nuclear Power Station



Analysis results of drainage water at Fukushima Daiichi Nuclear Power Station

Unit: Bq/L

	Drainage Channel A outlet							Drainage outlet at shallow draft quay						
Sampling date														
Sampling time														
Rainfall (mm/day)														
Flow rate (m3/minute)														
Cs-134 (Approx. 2 years)														
Cs-137 (Approx. 30 years)														
Gross β														
H-3 (Approx. 12 years)														

Unit: Bq/L

	Drainage Channel K outlet							Drainage Channel C at 35m above the sea level						
Sampling date														
Sampling time														
Rainfall (mm/day)														
Flow rate (m3/minute)														
Cs-134 (Approx. 2 years)														
Cs-137 (Approx. 30 years)														
Gross β														
H-3 (Approx. 12 years)														

• The latest figures are in the cells surrounded by thick lines.

• “-” indicates non-sampling target.

• “ND (not detected)” indicates that the measurement results are below the detection limits, and the detection limit of each radioactive material is provided in parentheses.

*The increase in values is presumed to have occurred because rainwater washed away surface soil, thereby carrying radioactive materials down to the drainage channel.

<Reference> The Highest Dose Until the Previous Measurement (Drainage water)

Unit: Bq/L

	Drainage Channel A outlet	Drainage outlet at shallow draft quay	Drainage Channel K outlet	Drainage Channel C at 35m above the sea level
Cs-134 (Approx. 2 years)				
Cs-137 (Approx. 30 years)				
Gross β				
H-3 (Approx. 12 years)				

* The sampling date is provided in parentheses.