

Nuclide Analysis Results of Fish and Shellfish
 <Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station>
 Samples collected in the third quarter of FY2019

[Measurement results of Sr-90 (half-life approx. 29 years) in fish]

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg(Raw)] (Half-life)	
			Sr-90 *1 (Approx. 29 years)	Reference*1 (Sum of Cs-134 and Cs-137)
Blue crab (whole) *2	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	0.078	19
Banded houndshark (muscle) *3	Around 2km Offshore of Kido River (T-S5)	November 6, 2019	0.010	5.6
Japanese angel shark (whole) *2	Around 2km Offshore of Kido River (T-S5)	December 20, 2019	0.023	19
Japanese angel shark (whole) *3	Around 2km Offshore of Fukushima Daini NPS(T-S7)	December 20, 2019	0.025	9.5
Blue crab (whole) *3	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	0.052	6.0
Japanese angel shark (whole) *2	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	0.019	14

*1 Edible parts (muscles) of fish were used to measure Cs. Whole fish (except for internal organs) including bones, which are not edible, were used to measure Sr.

Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*2 The Sr-90 analysis was conducted by KANSO CO., LTD.

*3 The Sr-90 analysis was conducted by Kyushu Environmental Evaluation Association.

Nuclide Analysis Results of Fish and Shellfish

<Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station>

Samples collected in the third quarter of FY2019

[Measurement results for Tritium (Half-life: Approx. 12 years) in fish and shellfish]

Place of Sampling(Place No.): Around 4km Offshore of Kumagawa (T-S8)

Name of Sample (Region)	Date of Sampling	Tritium concentration (Bq/L)		Tritium concentration (Bq/kg (Raw))		Reference (Sum of Cs-134 and Cs-137) (Bq/kg (Raw))
		Free Water Tritium	Organically Bound Tritium	Free Water Tritium	Organically Bound Tritium	
Flatfish ① (muscle)	October 31, 2019	0.094	ND(0.27)	0.075	ND(0.037)	ND
Flatfish ② (muscle)	November 21, 2019	0.073	ND(0.27)	0.058	ND(0.037)	ND
Flatfish (muscle)	December 13, 2019	0.083	ND(0.27)	0.065	ND(0.038)	ND

Reference

	Date of Sampling	Tritium concentration (Bq/L)
Around 4km Offshore of Kumagawa (T-S8) Seawater	October 30, 2019	0.10
	November 20, 2019	0.055
	December 12, 2019	0.078

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*The tritium analysis was conducted by Kyushu Environmental Evaluation Association.

*Edible parts (muscles) of fish were used to measure Cs.

*Free Water Tritium means tritium which is contained in the moisture of fish muscles and the values are compared with tritium concentrations in seawater where fish lives.

Organically Bound Tritium means tritium which is contained in dried fish muscles and the values show tritium concentrations in the vapor generated when dried fish is burned.

*The measurement results are rounded to two significant digits.

*ND indicates that a value is less than the detection limit of radioactive concentration. The detection limit is shown in parenthesis.