## Result of Sr nuclide analysis in the soil Fukushima Daiichi Nuclear Power Station

## 1. Measurement Result:

(Data summarized on April 18)

(Unit : Bq/kg dry soil)

Place of Sampling The Distance from Unit 1-2 Stacks in parentheses.	Date	Sr-89	Sr-90
(1) Ground (WNW approx. 500m)*1		N.D.	(6.5±0.12)×10 <sup>1</sup>
(2) Yachounomori (W approx. 500m)*1	Sep 16, 2013	N.D.	(1.2±0.017)×10 <sup>2</sup>
(3) Around industrial waste treatment facility (SSW approx.		N.D.	(1.1±0.016)×10 <sup>2</sup>
The range of the past measurement FY2008)*	results (FY1999 -	-	N.D 4.3

<sup>\*1</sup> Sampling was conducted in the area adjacent to the past sampling location to avoid duplication.

## 2. Analytical Institution: KAKEN Inc.

## 3. Evaluation:

The densities of Sr-90 are higher than those of the fallouts observed in Japan after the past atmospheric nuclear tests. Therefore, there is a possibility that the higher densities originate from the accident this time.

End

<sup>\*2</sup> Source "Report on the environmental radioactivity measurement around the Nuclear Power Plant (2008)", Committee on the safety technology of Nuclear Power Plants in Fukushima.