

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<1/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)           | Place of Sampling (Place No.)           | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|--------------------------------------|-----------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                      |                                         |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Greenling(muscle)                    | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(4.1)                                        | ND(3.5)                     | ND       |
| Northern dogfish(muscle)             | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(3.8)                                        | ND(3.6)                     | ND       |
| Blue crab(whole)                     | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(3.2)                                        | ND(4.0)                     | ND       |
| Schlegel's black<br>rockfish(muscle) | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(4.0)                                        | ND(3.6)                     | ND       |
| Sea raven(muscle)                    | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(4.0)                                        | ND(3.5)                     | ND       |
| Common skete(muscle)                 | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(4.0)                                        | 6.3                         | 6.3      |
| Sebastes cheni(muscle)               | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | 8.2                                            | 26                          | 34.2     |
| Drumfish(muscle)                     | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(3.0)                                        | ND(3.0)                     | ND       |
| Microstomus achne(muscle)            | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(4.1)                                        | 3.9                         | 3.9      |
| Ovalipes punctatus(whole)            | Around 1km Offshore of Ota River (T-S1) | June 11, 2015    | ND(3.6)                                        | ND(3.5)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<2/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)      | Place of Sampling (Place No.)            | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|---------------------------------|------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                 |                                          |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Flatfish(muscle)                | Around 1km Offshore of Ota River (T-S1)  | June 11, 2015    | ND(3.6)                                        | ND(3.5)                     | ND       |
| Northern dogfish(muscle)        | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.5)                                        | ND(3.0)                     | ND       |
| Stone flounder(muscle)          | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.7)                                        | ND(4.1)                     | ND       |
| Blue crab(whole)                | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.5)                                        | ND(3.5)                     | ND       |
| Lophius litilon(whole)          | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.0)                                        | ND(3.5)                     | ND       |
| Common skate(muscle)            | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(4.0)                                        | 18                          | 18       |
| Sebastes cheni(muscle)          | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | 9.0                                            | 43                          | 52       |
| Microstomus achne(muscle)       | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.3)                                        | 8.3                         | 8.3      |
| Flatfish(muscle)                | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.5)                                        | 3.8                         | 3.8      |
| Littlemouth<br>flounder(muscle) | Around 3km Offshore of Odaka Ward (T-S2) | June 11, 2015    | ND(3.7)                                        | ND(3.9)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<3/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)      | Place of Sampling (Place No.)             | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|---------------------------------|-------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                 |                                           |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Marbled sole(muscle)            | Around 3km Offshore of Odaka Ward (T-S2)  | June 11, 2015    | ND(3.5)                                        | ND(3.5)                     | ND       |
| Roundnose flounder(muscle)      | Around 3km Offshore of Odaka Ward (T-S2)  | June 11, 2015    | ND(3.5)                                        | ND(3.2)                     | ND       |
| Northern dogfish(muscle)        | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(3.2)                                        | ND(2.7)                     | ND       |
| Stone flounder(muscle)          | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(4.2)                                        | 12                          | 12       |
| Blue crab(whole)                | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(3.1)                                        | ND(4.0)                     | ND       |
| Lepidotrigla microptera(muscle) | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(3.8)                                        | ND(4.3)                     | ND       |
| Lophius litilon(whole)          | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(3.1)                                        | ND(3.6)                     | ND       |
| Common skete(muscle)            | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | 6.7                                            | 26                          | 32.7     |
| Starry flounder(muscle)         | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(4.5)                                        | 8.5                         | 8.5      |
| Ovalipes punctatus(whole)       | Around 3km Offshore of Ukedo River (T-S3) | June 10, 2015    | ND(4.2)                                        | ND(3.6)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg。

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<4/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)         | Place of Sampling (Place No.)                       | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|------------------------------------|-----------------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                    |                                                     |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Flatfish(muscle)                   | Around 3km Offshore of Ukedo River (T-S3)           | June 10, 2015    | ND(3.1)                                        | ND(3.8)                     | ND       |
| Marbled sole(muscle)               | Around 3km Offshore of Ukedo River (T-S3)           | June 10, 2015    | ND(3.4)                                        | 10                          | 10       |
| Northern dogfish(muscle)           | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.6)                                        | ND(3.6)                     | ND       |
| Blue crab(whole)                   | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.9)                                        | ND(4.5)                     | ND       |
| Lepidotrigla<br>microptera(muscle) | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.5)                                        | ND(3.5)                     | ND       |
| Lophius litilon(whole)             | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.5)                                        | ND(3.8)                     | ND       |
| Sea raven(muscle)                  | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.5)                                        | ND(3.6)                     | ND       |
| Common skete(muscle)               | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | 5.3                                            | 24                          | 29.3     |
| Banded dogfish(muscle)             | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.5)                                        | ND(3.6)                     | ND       |
| Microstomus achne(muscle)          | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | 6.3                                            | 28                          | 34.3     |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<5/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region) | Place of Sampling (Place No.)                       | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|----------------------------|-----------------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                            |                                                     |                  | Cs-134<br>(Approx. 2<br>years)                 | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Ovalipes punctatus(whole)  | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.3)                                        | ND(4.2)                     | ND       |
| Flatfish(muscle)           | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(4.0)                                        | ND(4.0)                     | ND       |
| Yellowtail(muscle)         | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.8)                                        | ND(3.7)                     | ND       |
| Marbled sole(muscle)       | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.7)                                        | 10                          | 10       |
| Chub mackerel(muscle)      | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.3)                                        | ND(3.4)                     | ND       |
| Roundnose flounder(muscle) | Around 3km Offshore of Fukushima Daiichi NPS (T-S4) | June 10, 2015    | ND(3.6)                                        | ND(3.6)                     | ND       |
| Greenling(muscle)          | Around 2km Offshore of Kido River (T-S5)            | June 5, 2015     | ND(3.8)                                        | 12                          | 12       |
| Common skete(muscle)       | Around 2km Offshore of Kido River (T-S5)            | June 5, 2015     | 6.3                                            | 26                          | 32.3     |
| Banded dogfish(muscle)     | Around 2km Offshore of Kido River (T-S5)            | June 5, 2015     | ND(4.7)                                        | ND(4.2)                     | ND       |
| Microstomus achne(muscle)  | Around 2km Offshore of Kido River (T-S5)            | June 5, 2015     | 4.7                                            | 20                          | 24.7     |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<6/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)           | Place of Sampling (Place No.)                         | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|--------------------------------------|-------------------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                      |                                                       |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Flatfish(muscle)                     | Around 2km Offshore of Kido River (T-S5)              | June 5, 2015     | ND(3.7)                                        | ND(3.6)                     | ND       |
| Smooth dogfish(muscle)               | Around 2km Offshore of Kido River (T-S5)              | June 5, 2015     | ND(3.3)                                        | ND(3.1)                     | ND       |
| Marbled sole(muscle)                 | Around 2km Offshore of Kido River (T-S5)              | June 5, 2015     | 4.5                                            | 19                          | 23.5     |
| Greenling(muscle)                    | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | ND(3.4)                                        | 9.9                         | 9.9      |
| Northern dogfish(muscle)             | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | ND(3.8)                                        | ND(3.2)                     | ND       |
| Schlegel's black<br>rockfish(muscle) | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | 8.8                                            | 31                          | 39.8     |
| Common skete(muscle)                 | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | 7.3                                            | 33                          | 40.3     |
| Banded dogfish(muscle)               | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | 10                                             | 37                          | 47       |
| Microstomus<br>achne(muscle)         | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | 19                                             | 65                          | 84       |
| Flatfish(muscle)                     | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | ND(4.2)                                        | ND(4.7)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<7/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)      | Place of Sampling (Place No.)                         | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                              |          |
|---------------------------------|-------------------------------------------------------|------------------|------------------------------------------------|------------------------------|----------|
|                                 |                                                       |                  | Cs-134<br>(Approx. 2 years)                    | Cs-137<br>(Approx. 30 years) | CS (Sum) |
| Dasyatis matsubarai(muscle)     | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | ND(3.0)                                        | ND(3.3)                      | ND       |
| Smooth dogfish(muscle)          | Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7) | June 5, 2015     | ND(3.7)                                        | 5.7                          | 5.7      |
| Northern dogfish(muscle)        | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.0)                                        | ND(3.3)                      | ND       |
| Blue crab(whole)                | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.9)                                        | ND(3.8)                      | ND       |
| Lepidotrigla microptera(muscle) | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.2)                                        | 5.1                          | 5.1      |
| Lophius litilon(whole)          | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.7)                                        | ND(3.6)                      | ND       |
| Common skete(muscle)            | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | 7.7                                            | 36                           | 43.7     |
| Ovalipes punctatus(whole)       | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(4.4)                                        | ND(3.6)                      | ND       |
| Flatfish(muscle)                | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.7)                                        | ND(4.5)                      | ND       |
| Littlemouth flounder(muscle)    | Around 4km Offshore of Kumagawa (T-S8)                | June 8, 2015     | ND(3.7)                                        | ND(3.5)                      | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<8/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)         | Place of Sampling (Place No.)             | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                              |          |
|------------------------------------|-------------------------------------------|------------------|------------------------------------------------|------------------------------|----------|
|                                    |                                           |                  | Cs-134<br>(Approx. 2<br>years)                 | Cs-137<br>(Approx. 30 years) | CS (Sum) |
| Marbled sole(muscle)               | Around 4km Offshore of Kumagawa (T-S8)    | June 8, 2015     | ND(4.2)                                        | 16                           | 16       |
| Greenling(muscle)                  | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(4.4)                                        | ND(3.5)                      | ND       |
| Stone flounder(muscle)             | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.9)                                        | 9.7                          | 9.7      |
| Lepidotrigla<br>microptera(muscle) | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(4.5)                                        | ND(3.8)                      | ND       |
| Lophius litilon(whole)             | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.5)                                        | ND(3.5)                      | ND       |
| Common skete(muscle)               | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.4)                                        | 10                           | 10       |
| Microstomus achne(muscle)          | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.7)                                        | 8.3                          | 8.3      |
| Flatfish(muscle)                   | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.6)                                        | 8.2                          | 8.2      |
| Littlemouth<br>flounder(muscle)    | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(4.1)                                        | ND(4.1)                      | ND       |
| Ridged-eye<br>flounder(muscle)     | Around 15km Offshore of Odaka Ward (T-B1) | June 12, 2015    | ND(3.9)                                        | ND(4.0)                      | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.



Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<9/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)      | Place of Sampling (Place No.)              | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                          |          |
|---------------------------------|--------------------------------------------|------------------|------------------------------------------------|--------------------------|----------|
|                                 |                                            |                  | Cs-134 (Approx. 2 years)                       | Cs-137(Approx. 30 years) | CS (Sum) |
| Greenling(muscle)               | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.2)                                        | ND(3.5)                  | ND       |
| Lepidotrigla microptera(muscle) | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.6)                                        | ND(3.8)                  | ND       |
| Lophius litilon(whole)          | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.3)                                        | ND(3.3)                  | ND       |
| Common skete(muscle)            | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(4.0)                                        | 13                       | 13       |
| Loliginid(whole)                | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.3)                                        | ND(4.1)                  | ND       |
| Cloudy catshark(muscle)         | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(4.4)                                        | 4.3                      | 4.3      |
| Microstomus achne(muscle)       | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(4.1)                                        | 4.7                      | 4.7      |
| Flatfish(muscle)                | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.8)                                        | ND(3.4)                  | ND       |
| Korean flounder(muscle)         | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.3)                                        | 3.7                      | 3.7      |
| Littlemouth flounder(muscle)    | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.5)                                        | ND(3.6)                  | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<10/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)         | Place of Sampling (Place No.)              | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|------------------------------------|--------------------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                    |                                            |                  | Cs-134<br>(Approx. 2<br>years)                 | Cs-137(Approx. 30<br>years) | CS (Sum) |
| Marbled sole(muscle)               | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.2)                                        | 8.6                         | 8.6      |
| Dory(muscle)                       | Around 18km Offshore of Ukedo River (T-B2) | June 12, 2015    | ND(3.9)                                        | ND(3.7)                     | ND       |
| Greenling(muscle)                  | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(3.9)                                        | ND(3.9)                     | ND       |
| Stone flounder(muscle)             | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | 4.5                                            | 18                          | 22.5     |
| Lepidotrigla<br>microptera(muscle) | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(3.5)                                        | ND(3.3)                     | ND       |
| Lophius litilon(whole)             | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(4.3)                                        | ND(3.3)                     | ND       |
| Common skete(muscle)               | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | 4.9                                            | 22                          | 26.9     |
| Microstomus achne(muscle)          | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(3.4)                                        | 7.8                         | 7.8      |
| Flatfish(muscle)                   | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(4.1)                                        | 6.1                         | 6.1      |
| Littlemouth<br>flounder(muscle)    | Around 10km Offshore of 1F (T-B3)          | June 16, 2015    | ND(3.1)                                        | ND(4.0)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)< 11/12  
>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)         | Place of Sampling (Place No.)     | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|------------------------------------|-----------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                                    |                                   |                  | Cs-134 (Approx.<br>2 years)                    | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Marbled sole(muscle)               | Around 10km Offshore of 1F (T-B3) | June 16, 2015    | ND(3.5)                                        | 13                          | 13       |
| Roundnose<br>flounder(muscle)      | Around 10km Offshore of 1F (T-B3) | June 16, 2015    | ND(4.3)                                        | ND(3.7)                     | ND       |
| Greenling(muscle)                  | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(3.5)                                        | ND(5.1)                     | ND       |
| Stone flounder(muscle)             | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(3.6)                                        | 3.7                         | 3.7      |
| Lepidotrigla<br>microptera(muscle) | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.0)                                        | ND(4.3)                     | ND       |
| Lophius litilon(whole)             | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.0)                                        | ND(4.0)                     | ND       |
| Common skete(muscle)               | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.2)                                        | 14                          | 14       |
| Microstomus achne(muscle)          | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(3.6)                                        | 11                          | 11       |
| Flatfish(muscle)                   | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.0)                                        | ND(3.4)                     | ND       |
| Littlemouth<br>flounder(muscle)    | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.2)                                        | ND(3.4)                     | ND       |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

\* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.

\* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS)<  
12/12>  
(excluding inside the port)

(Data summarized on July 17)

| Name of Sample<br>(Region)    | Place of Sampling (Place No.)     | Date of Sampling | Radioactivity Density[Bq/kg (Raw)] (Half-life) |                             |          |
|-------------------------------|-----------------------------------|------------------|------------------------------------------------|-----------------------------|----------|
|                               |                                   |                  | Cs-134<br>(Approx. 2<br>years)                 | Cs-137(Approx.<br>30 years) | CS (Sum) |
| Marbled sole(muscle)          | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.0)                                        | 13                          | 13       |
| Roundnose<br>flounder(muscle) | Around 10km Offshore of 2F (T-B4) | June 16, 2015    | ND(4.3)                                        | 5.3                         | 5.3      |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |
|                               |                                   |                  |                                                |                             |          |

\* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.  
 \* Standard value (since April 1, 2014) Total of Cs-134 and Cs-137=100Bq/kg.  
 \* Analyzed by: Tokyo Power Technology Ltd.