## Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station>

## (Data summarized on April 14)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Spotbelly rockfish (muscle)	Port area of Fukushima Daiichi NPS(Near south breakwater)	Mar. 10, 2017	8.1	48	56.1
Schlegel's black rockfish (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Mar. 23, 2017	12	85	97
Common skete (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Mar. 23, 2017	ND(8.9)	31	31
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Mar. 23, 2017	11	76	87
Barfin flounder (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Mar. 23, 2017	ND(5.7)	17	17

<sup>\*</sup>When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in paths as the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.