

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(4.0)	5.4	5.4
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.8)	ND(4.0)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.4)	8.9	8.9
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.6)	6.5	6.5
Sea bass (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.4)	ND(4.4)	ND
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.5)	4.6	4.6
Ovalipes punctaus (whole)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.3)	ND(4.3)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.6)	4.1	4.1
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.6)	4.7	4.7
Pacific cod (muscle)	Around 1km Offshore of Ota River (T-S1)	Apr. 14, 2017	ND(3.4)	3.9	3.9

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(3.3)	5.5	5.5
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(4.2)	ND(4.1)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(3.8)	ND(3.8)	ND
Sea bass (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(3.8)	ND(4.8)	ND
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(4.6)	ND(3.6)	ND
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(3.4)	3.8	3.8
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(4.0)	ND(4.5)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(3.9)	ND(3.7)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Apr. 14, 2017	ND(4.3)	ND(4.4)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.7)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.6)	ND(4.4)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.8)	ND(3.9)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.4)	5.9	5.9
Ovalipes punctaus (whole)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.8)	ND(4.0)	ND
Flat fish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.3)	ND(3.5)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.5)	ND(3.6)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(3.6)	6.3	6.3
Pacific cod (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Apr. 20, 2017	ND(4.2)	ND(3.3)	ND
Blue crab (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.0)	ND(4.0)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.9)	8.7	8.7

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(4.1)	9.4	9.4
Ovalipes punctatus (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.8)	ND(4.0)	ND
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.5)	ND(3.2)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.8)	ND(3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Apr. 20, 2017	ND(3.8)	ND(4.3)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.3)	4.6	4.6
Spiny dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.9)	3.9	3.9
Stone flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.9)	ND(4.2)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(4.0)	ND(3.5)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(4.0)	5.2	5.2

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.6)	16	16
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.7)	4.2	4.2
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.7)	5.2	5.2
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	Apr. 17, 2017	ND(3.8)	ND(3.9)	ND
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(3.7)	5.1	5.1
Stone flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(4.2)	ND(3.6)	ND
Common skate (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(4.2)	17	17
requiem shark (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(4.0)	ND(3.6)	ND
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(3.4)	13	13
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(3.4)	4.9	4.9

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Smooth dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(3.9)	ND(3.4)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Apr. 17, 2017	ND(3.6)	5.8	5.8
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.9)	ND(3.9)	ND
Lepidotrigla microptera (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.0)	ND(4.1)	ND
Schlegel's black rockfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.9)	8.2	8.2
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(4.0)	5.1	5.1
Pointhead flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.8)	ND(3.7)	ND
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(4.4)	15	15
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.6)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.5)	ND(4.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.0)	4.2	4.2
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Apr. 11, 2017	ND(3.4)	ND(3.9)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.7)	ND(4.1)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.5)	ND(3.7)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.5)	ND(3.9)	ND
Common skate (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(2.8)	ND(3.3)	ND
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.4)	ND(3.4)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.6)	ND(3.2)	ND
Flatfish① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(4.5)	ND(3.0)	ND
Flatfish② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(4.0)	ND(3.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.7)	ND(3.4)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(4.1)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.8)	ND(3.7)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(3.2)	8.6	8.6
Pacific cod (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Apr. 21, 2017	ND(4.2)	ND(3.3)	ND
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.6)	ND(3.8)	ND
golden cuttlefish (whole)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.4)	ND(3.9)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.9)	ND(3.3)	ND
Gray antimony(whole)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.1)	ND(3.7)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.9)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Dwarf squid (whole)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.5)	ND(4.2)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.9)	ND(3.2)	ND
Pointhead flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.6)	ND(2.5)	ND
Crimson sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.1)	ND(3.7)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.5)	ND(3.4)	ND
Flatfish① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.0)	ND(3.5)	ND
Flatfish② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.1)	ND(3.7)	ND
slime flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.2)	ND(3.5)	ND
Sea robin (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.3)	ND(4.3)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.9)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(4.0)	ND(4.1)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.6)	5.5	5.5
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.8)	ND(3.5)	ND
frog flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.5)	ND(3.8)	ND
Chestnut octopus(muscle)	Around 18km Offshore of Ukedo River (T-B2)	Apr. 21, 2017	ND(3.5)	ND(3.4)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.6)	ND(4.1)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.6)	ND(3.3)	ND
Schlegel's black rockfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(2.9)	ND(4.0)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.9)	6.2	6.2
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.6)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(4.5)	ND(3.8)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.3)	ND(3.4)	ND
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.6)	ND(3.9)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.2)	ND(3.7)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.3)	ND(3.9)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(4.1)	ND(4.1)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(4.2)	ND(3.4)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.9)	ND(3.8)	ND
Pacific cod (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(3.4)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND(5.0)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
frog flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Apr. 24, 2017	ND (3. 4)	ND (3. 7)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 4)	ND (4. 6)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 3)	ND (3. 0)	ND
Gray antimony(whole)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 9)	ND (3. 8)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (4. 1)	5. 1	5. 1
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 8)	ND (3. 0)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 5)	ND (3. 5)	ND
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (4. 0)	ND (3. 5)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (4. 1)	ND (4. 1)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 8)	ND (3. 4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <13/13>
(excluding the port)

(Data summarized on May 19)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (4. 2)	ND (4. 1)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (4. 0)	ND (3. 7)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 8)	ND (4. 2)	ND
frog flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Apr. 24, 2017	ND (3. 3)	ND (3. 7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.