

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stingray (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.1)	ND(3.9)	ND
Spiny dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.0)	ND(3.2)	ND
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.9)	ND(3.9)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.7)	5.8	5.8
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(4.1)	7.0	7.0
Drumfish (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.8)	ND(3.9)	ND
Ovalipes punctatus (whole)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.5)	ND(4.0)	ND
Flatfish① (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.9)	5.0	5.0
Flatfish② (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.9)	ND(3.3)	ND
Pitted stingray (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.7)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	June 15, 2017	ND(3.5)	5.6	5.6
Yellow goosfish (whole)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.1)	ND(3.8)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.5)	ND(3.6)	ND
Sea raven (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.0)	ND(3.6)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.1)	7.8	7.8
Banded houndshark (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.5)	ND(2.9)	ND
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.9)	ND(4.0)	ND
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.0)	ND(4.0)	ND
Flatfish① (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.8)	ND(3.7)	ND
Flatfish② (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.2)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(4.4)	ND(3.8)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.8)	ND(4.3)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.2)	3.4	3.4
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	June 15, 2017	ND(3.4)	ND(4.1)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.5)	ND(4.2)	ND
Lepidotrigla microptera (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.3)	ND(3.6)	ND
Yellow goosfish (whole)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.8)	ND(3.8)	ND
Microstomus achne (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.4)	ND(4.1)	ND
Flatfish① (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.3)	ND(3.6)	ND
Flatfish② (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(3.1)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(4.2)	ND(3.3)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	June 8, 2017	ND(4.5)	ND(4.5)	ND
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.6)	ND(4.2)	ND
Yellow goosfish (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.7)	ND(4.1)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.9)	7.8	7.8
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.5)	12	12
Flatfish① (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.8)	ND(4.2)	ND
Flatfish② (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(2.9)	ND(3.9)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(3.6)	ND(3.5)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	June 8, 2017	ND(4.0)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(4.3)	4.5	4.5
Stingray (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(3.5)	ND(3.9)	ND
Sea raven (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(4.1)	ND(3.8)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(3.1)	20	20
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(4.2)	6.6	6.6
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(3.7)	ND(4.1)	ND
Pitted stingray (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(3.9)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(3.3)	ND(4.0)	ND
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(4.1)	ND(3.3)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	June 6, 2017	ND(4.4)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(3.4)	ND(3.5)	ND
Fox jacopever (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(2.9)	6.3	6.3
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(3.7)	4.6	4.6
Banded houndshark (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(3.7)	7.2	7.2
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(4.4)	11	11
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(3.8)	4.2	4.2
Smooth dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(3.4)	5.5	5.5
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	June 6, 2017	ND(4.2)	ND(3.6)	ND
Lepidotrigla microptera (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.6)	ND(3.8)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(4.1)	6.0	6.0

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.8)	6.5	6.5
Flatfish① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.6)	ND(4.0)	ND
Flatfish② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(4.0)	ND(3.9)	ND
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(4.4)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.4)	ND(4.3)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.8)	5.5	5.5
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	June 15, 2017	ND(3.6)	ND(4.1)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(2.9)	ND(3.9)	ND
Striped jewfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(4.0)	ND(3.6)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.4)	ND(4.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Yellow goosfish (whole)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.8)	ND(4.1)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.6)	5.5	5.5
White croaker (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.5)	ND(3.1)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(4.0)	ND(3.9)	ND
Flatfish① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(4.5)	ND(4.5)	ND
Flatfish② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(4.2)	ND(4.0)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.8)	ND(4.6)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.7)	ND(3.1)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.9)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.1)	ND(3.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	June 2, 2017	ND(3.1)	ND(3.9)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.1)	ND(4.0)	ND
Yellow goosfish (whole)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(4.5)	ND(4.0)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.6)	ND(4.1)	ND
White croaker (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(4.3)	ND(3.9)	ND
Pointhead flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.9)	ND(4.2)	ND
Crimson sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.5)	ND(4.0)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.5)	ND(3.6)	ND
Flatfish① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.3)	ND(3.7)	ND
Flatfish② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(4.1)	3.7	3.7

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Blackfin flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(4.2)	ND(4.0)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.9)	ND(4.2)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.9)	ND(3.7)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.7)	6.6	6.6
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(4.1)	ND(2.9)	ND
Willow flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	June 2, 2017	ND(3.6)	ND(4.5)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.4)	ND(3.9)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(4.4)	ND(4.2)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.6)	ND(3.7)	ND
Yellow gosefish (whole)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.2)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.1)	7.9	7.9
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.3)	ND(3.9)	ND
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(4.4)	ND(3.4)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.4)	ND(3.9)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.5)	ND(3.4)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(4.1)	ND(4.0)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.9)	ND(4.2)	ND
John dory (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.7)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(4.1)	ND(4.1)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	June 26, 2017	ND(3.3)	ND(4.1)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(4.2)	ND(3.9)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.6)	ND(3.1)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.6)	ND(3.9)	ND
Yellow goosfish (whole)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.7)	ND(3.3)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.8)	5.1	5.1
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.0)	5.9	5.9
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.7)	ND(3.4)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.6)	ND(3.9)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.9)	ND(3.2)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.3)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <13/13>
(excluding the port)

(Data summarized on July 13)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.3)	ND(3.5)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	June 26, 2017	ND(3.8)	4.6	4.6

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.