## Result of Nuclide Analysis of Radioactive Materials in the Air at the Site of Fukushima Nuclear Power Stations

Reference

ata summarized on May 25)

Place of sampling	West gate of Fukushima Daiichi		MP-1 of Fukushima Daini (Reference)				
Date and time of sampling	11:20-11:50 May 24, 2011		9:12-9:22 May 24, 2011		15:06-15:16 May 24, 2011		②Density limit in the air to workers engaged in tasks associated with radiation (Bq/cm3) ※2
Detected nuclide (half-life)	①Radioactivity density※1 (Bq/cm3)	Scaling factor (1/2)	①Radioactivity density※1 (Bq/cm3)	Scaling factor (1)/2)	①Radioactivity density※1 (Bq/cm3)	Scaling factor (1)/2)	
I-131 (approx. 8 days)	2. 9E-06	0. 00	ND	-	ND	-	1E-03
Cs-134 (approx. 2 years)	8. 8E-06	0. 00	1. 3E-05	0. 01	1. 2E-05	0. 01	2E-03
Cs-137 (approx. 30 years)	1. 6E-05	0. 01	1. 2E-05	0.00	1. 4E-05	0.00	3E-03

<sup>31</sup> The value of Radioactivity density is the sum of the value of volatile nuclide's density and the value of paritulate nuclide's density

O.OE-O means  $O.O \times 10^{-O}$ 

Data of other nuclides are under examination.

X2 Statutory density limit to the 3-month average density of radioactive nuclides contained in the air that humans breathe.
In the case of more than 2 nuclides, summation of scaling factor for each statutory density is compared to 1.