Nuclide Analysis Results of Seawater < Offshore and Coast>

Attachment

(Data summarized on September 16)

Place of Sampling	North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel)		Channel of 1F		15 km offshore of Fukushima Daiichi Upper layer		15 km offshore of Fukushima Daini Upper layer		Density limit by the announcement of Reactor Regulation (Bq/L) (the density limit in the water outside of
Date of sampling	N/A		N/A		2011/8/27		2011/8/27		
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	areas in the section 6 of the appendix 2)
I-131 (about 8 days)	-	-	-	-	ND	-	ND	-	40
Cs-134 (about 2 years)	-	-	-	-	ND	-	ND	-	60
Cs-137 (about 30 years)	-	1	-	1	ND	-	ND	ı	90
Sr-89 (Approx. 51 days)	-	-	-	-	ND	-	ND	-	300
Sr-90 (Approx. 29 years)	-	-	-	-	ND	-	ND	-	30

^{*} Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L.

(Evaluation)

No Sr-89 or Sr-90 have been detected in the measured samples this time.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

^{*} I-131, Cs-134, Cs-137: released on August 28.

^{*} Analysis organization: Japan Chemical Analysis Center (Sr-89, Sr-90), TEPCO (I-131, Cs-134, Cs-137)