Nuclide Analysis Results of Seawater < Coast>

Reference

(Data summarized on December 5)

Place of Sampling	North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel)		Around South Discharge Channel of 1F (appox. 330m south of 1-4u Discharge Channel)		Around North Discharge Channel of 2F (Around 3,4u Discharge Channel) (approx. 10 km from 1F)		Around Iwasawa Shore of 2F (appox. 7 km south of 1,2u Discharge Channel) (appox. 16 km from 1F)		Doncity limit by the
Time of Sampling	Dec 04, 2011 (Not sampled)		Dec 04, 2011 (Not sampled)		Dec 04, 2011 (Not sampled)		Dec 04, 2011 08:00 am		
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	areas in the section 6 of the appendix 2)
I-131 (about 8 days)	-	-	-	-	-	-	ND	-	40
Cs-134 (about 2 years)	-	-	-	-	1	-	ND	-	60
Cs-137 (about 30 years)	-	-	-	-	-	-	1.2	0.01	90

^{*} Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L.

I-131: approx. 0.66Bq/L, Cs-134: approx. 0.89Bq/L

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

^{*} Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

^{* &}quot;ND" means the sampled data is below measurable limit.